

# CONCUSSION FACT SHEET FOR PARENTS



## WHAT IS A CONCUSSION?

A concussion is a type of traumatic brain injury. Concussions are caused by a bump or blow to the head. Even a “ding,” “getting your bell rung,” or what seems to be a mild bump or blow to the head can be serious.

You can’t see a concussion. Signs and symptoms of concussion can show up right after the injury or may not appear or be noticed until days or weeks after the injury. If your child reports any symptoms of concussion, or if you notice the symptoms yourself, seek medical attention right away.

## WHAT ARE THE SIGNS AND SYMPTOMS OF CONCUSSION?

If your child has experienced a bump or blow to the head during a game or practice, look for any of the following signs of a concussion:

### SYMPTOMS REPORTED BY ATHLETE:

- Headache or “pressure” in head
- Nausea or vomiting
- Balance problems or dizziness
- Double or blurry vision
- Sensitivity to light
- Sensitivity to noise
- Feeling sluggish, hazy, foggy, or groggy
- Concentration or memory problems
- Confusion
- Just not “feeling right” or is “feeling down”

### SIGNS OBSERVED BY PARENTS/ GUARDIANS:

- Appears dazed or stunned
- Is confused about assignment or position
- Forgets an instruction
- Is unsure of game, score, or opponent
- Moves clumsily
- Answers questions slowly
- Loses consciousness (even briefly)
- Shows mood, behavior, or personality changes



## DANGER SIGNS

Be alert for symptoms that worsen over time. Your child or teen should be seen in an emergency department right away if s/he has:

- One pupil (the black part in the middle of the eye) larger than the other
- Drowsiness or cannot be awakened
- A headache that gets worse and does not go away
- Weakness, numbness, or decreased coordination
- Repeated vomiting or nausea
- Slurred speech
- Convulsions or seizures
- Difficulty recognizing people or places
- Increasing confusion, restlessness, or agitation
- Unusual behavior
- Loss of consciousness (even a brief loss of consciousness should be taken seriously)

## WHAT SHOULD YOU DO IF YOU THINK YOUR CHILD HAS A CONCUSSION?

### 1. SEEK MEDICAL ATTENTION RIGHT AWAY

A health care professional will be able to decide how serious the concussion is and when it is safe for your child to return to regular activities, including sports.

### 2. KEEP YOUR CHILD OUT OF PLAY.

Concussions take time to heal. Don't let your child return to play the day of the injury and until a health care professional says it's OK. Children who return to play too soon - while the brain is still healing - risk a greater chance of having a second concussion. Repeat or later concussions can be very serious. They can cause permanent brain damage, affecting your child for a lifetime.

### 3. TELL YOUR CHILD'S COACH ABOUT ANY PREVIOUS CONCUSSION.

Coaches should know if your child had a previous concussion. Your child's coach may not know about a concussion your child received in another sport or activity unless you tell the coach.

## HOW CAN YOU HELP YOUR CHILD PREVENT A CONCUSSION OR OTHER SERIOUS BRAIN INJURY?

- Ensure that they follow their coach's rules for safety and the rules of the sport.
- Encourage them to practice good sportsmanship at all times.
- Make sure they wear the right protective equipment for their activity. Protective equipment should fit properly and be well maintained.
- Wearing a helmet is a must to reduce the risk of a serious brain injury or skull fracture.
  - However, helmets are not designed to prevent concussions. There is no "concussion-proof" helmet. So, even with a helmet, it is important for kids and teens to avoid hits to the head.

## HOW CAN I HELP MY CHILD RETURN TO SCHOOL SAFELY AFTER A CONCUSSION?

Children and teens who return to school after a concussion may need to:

- Take rest breaks as needed
- Spend fewer hours at school
- Be given more time to take tests or complete assignments
- Receive help with schoolwork
- Reduce time spent reading, writing, or on the computer

Talk with your child's teachers, school nurse, coach, speech-language pathologist, or counselor about your child's concussion and symptoms. As your child's symptoms decrease, the extra help or support can be removed gradually.



JOIN THE CONVERSATION  [www.facebook.com/CDCHeadsUp](http://www.facebook.com/CDCHeadsUp)

TO LEARN MORE GO TO >> [WWW.CDC.GOV/CONCUSSION](http://WWW.CDC.GOV/CONCUSSION)

Content Source: CDC's Heads Up Program. Created through a grant to the CDC Foundation from the National Operating Committee on Standards for Athletic Equipment (NOCSAE).



## HEADS UP CONCUSSION

### Benjamin Franklin High School

## FACT SHEET FOR HIGH SCHOOL ATHLETES

This sheet has information to help you protect yourself from concussion or other serious brain injury and know what to do if a concussion occurs.

### WHAT IS A CONCUSSION?

---

A concussion is a brain injury that affects how your brain works. It can happen when your brain gets bounced around in your skull after a fall or hit to the head.

### WHAT SHOULD I DO IF I THINK I HAVE A CONCUSSION?

---

#### REPORT IT.

Tell your coach, parent, and athletic trainer if you think you or one of your teammates may have a concussion. It's up to you to report your symptoms. Your coach and team are relying on you. Plus, you won't play your best if you are not feeling well.

#### GET CHECKED OUT.

If you think you have a concussion, do not return to play on the day of the injury. Only a health care provider can tell if you have a concussion and when it is OK to return to school and play. The sooner you get checked out, the sooner you may be able to safely return to play.

#### GIVE YOUR BRAIN TIME TO HEAL.

A concussion can make everyday activities, such as going to school, harder. You may need extra help getting back to your normal activities. Be sure to update your parents and doctor about how you are feeling.



## WHY SHOULD I TELL MY COACH AND PARENT ABOUT MY SYMPTOMS?

- Playing or practicing with a concussion is dangerous and can lead to a longer recovery.
- While your brain is still healing, you are much more likely to have another concussion. This can put you at risk for a more serious injury to your brain and can even be fatal.

## HOW CAN I TELL IF I HAVE A CONCUSSION?

You may have a concussion if you have any of these symptoms after a bump, blow, or jolt to the head or body:

- Get a headache
- Feel dizzy, sluggish, or foggy
- Are bothered by light or noise
- Have double or blurry vision
- Vomit or feel sick to your stomach
- Have trouble focusing or problems remembering
- Feel more emotional or "down"
- Feel confused
- Have problems with sleep

Concussion symptoms usually show up right away, but you might not notice that something "isn't right" for hours or days. A concussion feels different to each person, so it is important to tell your parents and doctor how you are feeling.

The information provided in this document or through linkages to other sites is not a substitute for medical or professional care. Questions about diagnosis and treatment for concussion should be directed to a physician or other health care provider.

## HOW CAN I HELP MY TEAM?

### PROTECT YOUR BRAIN.

Avoid hits to the head and follow the rules for safe and fair play to lower your chances of getting a concussion. Ask your coaches for more tips.

### BE A TEAM PLAYER.

You play an important role as part of a team. Encourage your teammates to report their symptoms and help them feel comfortable taking the time they need to get better.

### GOOD TEAMMATES KNOW:

**"IT'S BETTER TO MISS ONE GAME THAN THE WHOLE SEASON."**



TO LEARN MORE GO TO

>> [www.cdc.gov/HEADSUP](http://www.cdc.gov/HEADSUP)

JOIN THE CONVERSATION AT

↳ [www.facebook.com/CDCHEADSUP](http://www.facebook.com/CDCHEADSUP)



Content Source: CDC's HEADS UP campaign. Customizable HEADS UP fact sheets were made possible through a grant to the CDC Foundation from the National Operating Committee on Standards for Athletic Equipment (NOCSAE).